

SUPERGRAVITY UNIFIED MODELS

R. ARNOWITT^a and PRAN NATH^b^a Center for Theoretical Physics, Department of Physics, Texas A&M University,
College Station, TX 77843-4242^b Department of Physics, Northeastern University, Boston, MA 02115**Abstract**

The development of supergravity unified models and their implications for current and future experiment are discussed.

1. Introduction

Supersymmetry (SUSY) was initially introduced as a global symmetry¹ on purely theoretical grounds that nature should be symmetric between bosons and fermions. It was soon discovered, however, that models of this type had a number of remarkable properties². Thus the bose-fermi symmetry led to the cancellation of a number of the infinities of conventional field theories, in particular the quadratic divergences in the scalar Higgs sector of the Standard Model (SM). Thus SUSY could resolve the gauge hierarchy problem that plagued the SM. Further, the hierarchy problem associated with grand unified models (GUT)³, where without SUSY, loop corrections gave all particles GUT size masses⁴ was also resolved. In addition, SUSY GUT models with minimal particle spectrum raised the value for the scale of grand unification, M_G , to $M_G \cong 2 \times 10^{16}$ GeV, so that the predicted proton decay rate^{5,6} was consistent with existing experimental bounds. Thus in spite of the lack of any direct experimental evidence for the existence of SUSY particles, supersymmetry became a highly active field among particle theorists.

However, by about 1980, it became apparent that global supersymmetry was unsatisfactory in that a phenomenologically acceptable picture of spontaneous breaking of supersymmetry did not exist. Thus the success of the SUSY grand unification program discussed above was in a sense spurious in that the needed SUSY threshold M_S (below which the SM held) could not be theoretically constructed. In order to get a phenomenologically viable model, one needed “soft breaking” masses (i.e. supersymmetry breaking terms of dimension ≤ 3 which maintain the gauge hierarchy⁷) and these had to be

introduced in an ad hoc fashion by hand. In the minimal SUSY model, the MSSM⁸, where the particle spectrum is just that of the supersymmetrized SM, one could introduce as many as 105 additional parameters (62 new masses and mixing angles and 43 new phases) leaving one with a theory with little predictive power.

A resolution of the problem of how to break supersymmetry spontaneously was achieved by promoting supersymmetry to a local symmetry i.e. supergravity⁹. Here gravity is included into the dynamics. One can then construct supergravity [SUGRA] grand unified models^{10,11} where the spontaneous breaking of supersymmetry occurs in a “hidden” sector via supergravity interactions in a fashion that maintains the gauge hierarchy. In such theories there remains, however, the question of at what scale does supersymmetry break, and what is the “messenger” that communicates this breaking from the hidden to the physical sector. In this chapter we consider models where supersymmetry breaks at a scale $Q > M_G$ with gravity being the messenger¹². Such models are economical in that both the messenger field and the agency of supersymmetry breaking are supersymmetrized versions of fields and interactions that already exist in nature (i.e. gravity). In addition, models of this type may turn out to be consequences of string theory.

The strongest direct evidence supporting supergravity GUT models is the apparent experimental grand unification of the three gauge coupling constants¹³. This result is non-trivial not only because three lines do not ordinarily intersect at one point, but also because there is only a narrow acceptable window for M_G . Thus one requires $M_G \gtrsim 5 \times 10^{15} \text{ GeV}$ so as not to violate current experimental bounds on proton decay for the $p \rightarrow e^+ + \pi^0$ channel (which occurs in almost all GUT models) and one requires $M_G \lesssim 5 \times 10^{17} \text{ GeV} \cong M_{string}$ (the string scale) so that gravitational effects do not become large invalidating the analysis. Further, assuming an MSSM type of particle spectrum between the electroweak scale M_Z and M_G , acceptable grand unification occurs only with one pair of Higgs doublets and at most four generations. Finally, naturalness requires that SUSY thresholds be at $M_S \lesssim 1 \text{ TeV}$ which turns out to be the case. Thus the possibility of grand unification is tightly constrained.

At present, grand unification in SUGRA GUTs can be obtained to within about 2-3 std.^{14,15} However, the closeness of M_G to the Planck scale, $M_{Pl} = (\hbar c / 8\pi G_N)^{1/2} \cong 2.4 \times 10^{18} \text{ GeV}$, suggests the possibility that there are $O(M_G/M_{Pl})$ corrections to these models. One might, in fact, expect such structures to arise in string theory as nonrenormalizable operators (NROs) obtained upon integrating out the tower of Planck mass states. Such terms would produce $\approx 1\%$ corrections at M_G which might grow to $\approx 5\%$ corrections at M_Z . Indeed, as will be seen in Sec.2, it is just such NRO terms involving

the hidden sector fields that give rise to the soft breaking masses, and so it would not be surprising to find such structures in the physical sector as well. Thus SUGRA GUTs should be viewed as an effective theory and, as will be discussed in Sec.4, with small deviations between theory and experiment perhaps opening a window to Planck scale physics.

One of the fundamental aspects of the SM, not explained by that theory, is the origin of the spontaneous breaking of $SU(2) \times U(1)$. SUGRA GUTs offers an explanation of this due to the existence of soft SUSY breaking masses at M_G . Thus as long as at least one of the soft breaking terms are present at M_G , breaking of $SU(2) \times U(1)$ can occur at a lower energy^{10,16}, providing a natural Higgs mechanism. Further, radiative breaking occurs at the electroweak scale provided the top quark is heavy ie. $100 \text{ GeV} \lesssim m_t \lesssim 200 \text{ GeV}$. The minimal SUGRA model^{10,17-19} (MSGM), which assumes universal soft breaking terms, requires only four additional parameters and one sign to describe all the interactions and masses of the 32 SUSY particles. Thus the MSGM is a reasonably predictive model, and for that reason is the model used in much of the phenomenological analysis of the past decade. However, we will see in Sec. 2 that there are reasons to consider non-universal extensions of the MSGM, and discuss some of the experimental consequences they produce in Sec.4.

2. Soft Breaking Masses

Supergravity interactions with chiral matter fields, $\{\chi_i(x), \phi_i(x)\}$ (where $\chi_i(x)$ are left (L) Weyl spinors and $\phi_i(x)$ are complex scalar fields) depend upon three functions of the scalar fields: the superpotential $W(\phi_i)$, the gauge kinetic function $f_{\alpha\beta}(\phi_i, \phi_i^\dagger)$ (which enters in the Lagrangian as $f_{\alpha\beta} F_{\mu\nu}^\alpha F^{\mu\nu\beta}$ with α, β = gauge indices) and the Kahler potential $K(\phi_i, \phi_i^\dagger)$ (which appears in the scalar kinetic energy as $K_j^i \partial_\mu \phi_i \partial^\mu \phi_j^\dagger$, $K_j^i \equiv \partial^2 K^2 / \partial \phi_i \partial \phi_j^\dagger$ and elsewhere). W and K enter only in the combination

$$G(\phi_i, \phi_i^\dagger) = \kappa^2 K(\phi_i, \phi_i^\dagger) + \ell n[\kappa^6 |W(\phi_i)|^2] \quad (1)$$

where $\kappa = 1/M_{Pl}$. Writing $\{\phi_i\} = \{\phi_a, z\}$ where ϕ_a are physical sector fields (squarks, sleptons, higgs) and z are the hidden sector fields whose VEVs $\langle z \rangle = \mathcal{O}(M_{Pl})$ break supersymmetry, one assumes that the superpotential decomposes into a physical and a hidden part,

$$W(\phi_i) = W_{phy}(\phi_a, \kappa z) + W_{hid}(z) \quad (2)$$

Supersymmetry breaking is scaled by requiring $\kappa^2 W_{hid} = \mathcal{O}(M_S) \tilde{W}_{hid}(\kappa z)$ and the gauge hierarchy is then guaranteed by the additive nature of the terms in

Eq.(2). Thus only gravitational interactions remain to transmit SUSY breaking from the hidden sector to the physical sector.

A priori, the functions W , K and $f_{\alpha\beta}$ are arbitrary. However, they are greatly constrained by the conditions that the model correctly reduce to the SM at low energies, and that non-renormalizable corrections be scaled by κ (as would be expected if they were the low energy residue of string physics of the Planck scale). Thus one can expand these functions in polynomials of the physical fields ϕ_a

$$f_{\alpha\beta}(\phi_i, \phi_i^\dagger) = c_{\alpha\beta} + \kappa d_{\alpha\beta}^a(x, y) \phi_a + \dots \quad (3)$$

$$W_{phys}(\phi_i) = \frac{1}{6} \lambda^{abc}(x) \phi_a \phi_b \phi_c + \frac{1}{24} \kappa \lambda^{abcd}(x) \phi_a \phi_b \phi_c \phi_d + \dots \quad (4)$$

$$\begin{aligned} K(\phi_i, \phi_i^\dagger) &= \kappa^{-2} c(x, y) + c_b^a(x, y) \phi_a \phi_b^\dagger + (c^{ab}(x, y) \phi_a \phi_b + h.c.) \\ &+ \kappa (c_{bc}^a \phi_a \phi_b^\dagger \phi_c^\dagger + h.c.) + \dots \end{aligned} \quad (5)$$

$x = \kappa z$ and $y = \kappa z^\dagger$, so that $\langle x \rangle, \langle y \rangle = \mathcal{O}(1)$. The scaling hypothesis for the NRO's imply then that the VEVs of the coefficients $c_{\alpha\beta}$, $c_{\alpha\beta}^a$, λ^{abc} , c , c_b^a , c^{ab} etc. are all $\mathcal{O}(1)$.

The holomorphic terms in K labeled by c^{ab} can be transformed from K to W by a Kahler transformation, $K \rightarrow K - (c^{ab} \phi_a \phi_b + h.c.)$ and

$$W \rightarrow W \exp[\kappa^2 c^{ab} \phi_a \phi_b] = W + \tilde{\mu}^{ab} \phi_a \phi_b + \dots \quad (6)$$

where $\tilde{\mu}^{ab}(x, y) = \kappa^2 W c^{ab}$. Hence $\langle \tilde{\mu}^{ab} \rangle = \mathcal{O}(\mathcal{M}_S)$, and one obtains a μ -term with the right order of magnitude after SUSY breaking provided only that c^{ab} is not zero²⁰. The cubic terms in W are just the Yukawa couplings with $\langle \lambda^{abc}(x) \rangle$ being the Yukawa coupling constants. Also $\langle c_{\alpha\beta} \rangle = \delta_{\alpha\beta}$, $\langle c_b^a \rangle = \delta_b^a$ and $\langle c_{xy} \rangle = 1$ ($c_x \equiv \partial c / \partial x$ etc.) so that the field kinetic energies have canonical normalization.

The breaking of SUSY in the hidden sector leads to the generation of a series of soft breaking terms^{10,11,17-21}. We consider here the case where $\langle x \rangle = \langle y \rangle$ (i.e. the hidden sector SUSY breaking does not generate any CP violation) and state the leading terms. Gauginos gain a soft breaking mass term at M_G of $(m_{1/2})_{\alpha\beta} \lambda^\alpha \gamma^0 \lambda^\beta$ (λ^α = gaugino Majorana field) where

$$(m_{1/2})_{\alpha\beta} = \kappa^{-2} \langle G^i (K^{-1})_j^i \text{Re} f_{\alpha\beta j}^\dagger \rangle m_{3/2} \quad (7)$$

Here $G^i \equiv \partial G / \partial \phi_i$, $(K^{-1})_j^i$ is the matrix universe of K_j^i , $f_{\alpha\beta j} = \partial f_{\alpha\beta} / \partial \phi_j^\dagger$ and $m_{3/2}$ is the gravitino mass: $m_{3/2} = \kappa^{-1} \langle \exp[G/2] \rangle$. In terms of the expansion of Eqs.(3-5) one finds

$$(m_{1/2})_{\alpha\beta} = [c + \ell n(W_{hid})]_x \text{Re} c_{\alpha\beta y}^* m_{3/2} \quad (8)$$

and $m_{3/2} = (\exp \frac{1}{2}c)\kappa^2 W_{hid}$ where it is understood from now on that x is to be replaced by its VEV in all functions (e.g. $c(x) \rightarrow c(\langle x \rangle) = \mathcal{O}(1)$) so that $m_{3/2} = \mathcal{O}(M_S)$. One notes the following about Eq.(8): (i) For a simple GUT group, gauge invariance implies that $c_{\alpha\beta} \sim \delta_{\alpha\beta}$ and so gaugino masses are universal at mass scales above M_G . (ii) While $m_{1/2}$ is scaled by $m_{3/2} = \mathcal{O}(M_S)$, it can differ from it by a significant amount. (iii) From Eq.(8) one sees that it is the NRO such as $\kappa z m_{3/2} \lambda^\alpha \gamma^0 \lambda^\alpha$ that gives rise to $m_{1/2}$. Below M_G , where the GUT group is broken, the second term of Eq.(3) would also contribute yielding a NRO of size $\kappa d_{\alpha\beta}^a \phi_a m_{3/2} \lambda^\alpha \gamma^0 \lambda^\beta \sim (M_G/M_{Pl}) m_{3/2} \lambda \gamma^0 \lambda$ ²³ for fields with VEV $\langle \phi_a \rangle = \mathcal{O}(M_G)$ which break the GUT group. Such terms give small corrections to the universality of the gaugino masses and effect grand unification. They are discussed in Sec.4.

The effective potential for the scalar components of chiral multiplets is given by ^{10,22}

$$V = \exp[\kappa^2 K] [(K^{-1})_i^j (W^i + \kappa^2 K^i W) (W^j + \kappa^2 K^j W)^\dagger - 3\kappa^2 |W|^2] + V_D \quad (9)$$

where $W^i = \partial W / \partial \phi_i$ etc., and

$$V_D = \frac{1}{2} g_\alpha g_\beta (Ref^{-1})_{\alpha\beta} (K^i (T^\alpha)_{ij} \phi_j) (K^k (T^\beta)_{kl} \phi_l) \quad (10)$$

where g_α are the gauge coupling constants. Eqs.(2-6) then lead to the following soft breaking terms at M_G :

$$V_{soft} = (m_0^2)_b^a \phi_a \phi_b^\dagger + \left[\frac{1}{3} \tilde{A}^{abc} \phi_a \phi_b \phi_c + \frac{1}{2} \tilde{B}^{ab} \phi_a \phi_b + h.c. \right] \quad (11)$$

In the following, we impose the condition that the cosmological constant vanish after SUSY breaking, i.e. $\langle V \rangle = 0$. [This is a fine tuning of $\mathcal{O}(M_S^2 M_{Pl}^2)$.] Then one finds for D-flat breaking

$$(m_0^2)_b^a = [3(c_{cx}^a c_{by}^c - c_b^a c_{xy}) + \delta_b^a] m_{3/2}^2 \quad (12)$$

$$\tilde{A}^{abc} = \pm \sqrt{3} \left[\left\{ c_x h^{abc} - (c_{dx}^a h^{dac} + c_{dx}^b h^{dac} + c_{dx}^c h^{abd}) \right\} + h_x^{abc} \right] m_{3/2} \quad (13)$$

$$\tilde{B}^{ab} = \left[\left\{ (\pm \sqrt{3} c_x - 1) \mu^{ab} \mp \sqrt{3} (c_{dx}^a \mu^{db} + c_{dx}^b \mu^{ad}) \right\} \pm \sqrt{3} \mu_x^{ab} \right] m_{3/2} \quad (14)$$

where the sign ambiguity arises from the two roots of the equation $\langle V \rangle = 0$. In Eqs.(12-14), $h^{abc} = \langle \exp(\frac{1}{2}c) \lambda^{abc} \rangle$ are the Yukawa coupling constants and $\mu^{ab} = \langle \exp(\frac{1}{2}c) \tilde{\mu}^{ab} \rangle$ is the μ -parameter, $h_x^{abc} \equiv \langle \exp(\frac{1}{2}c) \partial \lambda^{abc} / \partial x \rangle$ and $\mu_x^{ab} \equiv \langle \exp(\frac{1}{2}c) \partial \tilde{\mu}^{ab} / \partial x \rangle$. Since K is hermitian, c_b^a (defined in Eq.(5)) obeys $c_b^a = c_a^{b*}$.

One notes the following about Eqs.(12-14): (i) The scalar soft breaking masses $(m_0^2)_b^a$ are in general not universal unless the Kahler metric is flat, i.e.

$c_b^a = \delta_b^a$ ^{10,20}, or more generally unless the fields z which break SUSY couple universally to the physical sector i.e. $c_b^a = \tilde{c}(x, y)\delta_b^a$. (ii) $\tilde{A}^{abc}$ can contain a term, λ_x^{abc} which is not scaled by the Yukawa coupling constants h^{abc} if λ^{abc} is a function of the z fields. This possibility may occur in string theory, where the role of z is possibly played by the dilaton and moduli fields²¹. In the following we will neglect the λ_x^{abc} term (and similarly the μ_x^{ab} term in Eq.(14)).

The soft breaking masses in Eqs.(12-14) depend upon two hermitian matrices c_{bx}^a and c_{bxy}^a and the parameter c_x . The number of new SUSY parameters this implies depends upon the GUT group. We consider here the example of SU(5) supergravity GUT with R parity invariance. The superpotential is given by

$$W_{phys} = h_{ij}^{(1)} M_i^{XY} \overline{M}_{Xj} H_{1Y} + \frac{1}{4} h_{ij}^{(2)} \epsilon_{XYZWU} M_i^{XY} M_j^{ZW} H_2^U + \mu H_2^X \overline{H}_{1X} + W_{GUT} \quad (15)$$

where $X, Y, \dots = 1 \dots 5$ are SU(5) indices, $i, j = 1, 2, 3$ are generation indices, M_i^{XY} and $\overline{M}_{Xj}$ are 10 and $\overline{5}$ matter fields and H_2^X and $\overline{H}_{1X}$ are 5 and $\overline{5}$ Higgs fields. W_{GUT} contains additional Higgs fields which break SU(5) at M_G . H_2 and $\overline{H}_1$ contain the pair of light Higgs doublets below M_G . The soft breaking parameters then lead to 42 additional parameters for this model (25 new masses and mixing angles and 17 new phases), considerably fewer than the 105 additional parameters of the MSSM. While this is still a formidable number, many are experimentally uninteresting (e.g. scaled by the small Yukawa coupling constants), and we will see in Sec.4 that the remaining parameters are few enough (at least for the CP conserving sectors of the theory) to obtain interesting experimental predictions.

3. Radiative Breaking and the Low Energy Theory

In Sec.2, the SUGRA GUT model above the GUT scale i.e. at $Q > M_G$ was discussed. Below M_G the GUT group is spontaneously broken, and we will assume here that the SM group, SU(3) x SU(2) x U(1), holds for $Q < M_G$ ²⁴. Contact with accelerator physics at low energy can then be achieved using the renormalization group equations (RGE) running from M_G to the electroweak scale M_Z . As one proceeds downward from M_G , the coupling constants and masses evolve, and provided at least one soft breaking parameter and also the μ parameter at M_G is not zero, the large top quark Yukawa can turn the H_2 running (mass)², $m_{H_2}^2(Q)$, negative at the electroweak scale¹⁶. Thus the spontaneous breaking of supersymmetry at M_G triggers the spontaneous

breaking of $SU(2) \times U(1)$ at the electroweak scale. In this fashion all the masses and coupling constants at the electroweak scale can be determined in terms of the fundamental parameters (Yukawa coupling constants and soft breaking parameters) at the GUT scale, and the theory can be subjected to experimental tests.

The conditions for electroweak symmetry breaking arise from minimizing the effective potential V at the electroweak scale with respect to the Higgs VEVs $v_{1,2} = \langle H_{1,2} \rangle$. This leads to the equations¹⁶

$$\mu^2 = \frac{\mu_1^2 - \mu_2^2 \tan^2 \beta}{\tan^2 \beta - 1} - \frac{1}{2} M_Z^2; \quad \sin^2 \beta = \frac{-2B\mu}{2\mu^2 + \mu_1^2 + \mu_2^2} \quad (16)$$

where $\tan \beta = v_2/v_1$, B is the quadratic soft breaking parameter ($V_{soft}^B = B\mu H_1 H_2$), $\mu_i = m_{H_i}^2 + \Sigma_i$, and Σ_i are loop corrections²⁶. All parameters are running parameters at the electroweak scale which we take for convenience to be $Q = M_Z$. Eq.(16) then determines the μ parameter and allows the elimination of B in terms of $\tan \beta$. This determination of μ greatly enhances the predictive power of the model.

We parameterize the Higgs soft breaking masses at M_G by

$$m_{H_1}^2 = m_0^2(1 + \delta_1); \quad m_{H_2}^2 = m_0^2(1 + \delta_2) \quad (17)$$

and the third generation of sfermion masses by

$$\begin{aligned} m_{q_L}^2 &= m_0^2(1 + \delta_3); \quad m_{u_R}^2 = m_0^2(1 + \delta_4); \quad m_{e_R}^2 = m_0^2(1 + \delta_5) \\ m_{d_R}^2 &= m_0^2(1 + \delta_6); \quad m_{\ell_L}^2 = m_0^2(1 + \delta_7); \end{aligned} \quad (18)$$

where $q_L = (u_L, d_L)$, $\ell_L = (\nu_L, e_L)$ etc. The δ_i thus represent non-universal deviations from the reference mass m_0 . For GUT groups which contain an $SU(5)$ subgroup (e.g. $SU(N)$, $N \geq 5$; $SO(N)$, $N \geq 10$, E_6 etc.) with matter embedded in the 10 and $\bar{5}$ representations, one has

$$\delta_3 = \delta_4 = \delta_5; \quad \delta_6 = \delta_7 \quad (19)$$

In the following we will choose m_0 to be the soft breaking mass of the first two generations (assumed universal to suppress FCNC effects⁵). We will also keep only the large top Yukawa coupling constant h_t and its associated cubic soft SUSY breaking parameter A_t (an approximation valid for $\tan \beta \lesssim 20$). This allows an analytic solution of the 1-loop RGE and enables one to understand physically the predictions of the model. One finds²⁷

$$\begin{aligned} \mu^2 &= \frac{t^2}{t^2 - 1} \left[\left(\frac{1 - 3D_0}{2} + \frac{1}{t^2} \right) + \left(\frac{1 - D_0}{2} (\delta_3 + \delta_4) - \frac{1 + D_0}{2} \delta_2 \right. \right. \\ &\quad \left. \left. + \frac{1}{t^2} \delta_1 \right) \right] m_0^2 + \frac{t^2}{t^2 - 1} \left[\frac{1}{2} (1 - D_0) \frac{A_R^2}{D_0} + C_\mu m_{1/2}^2 \right] - \frac{1}{2} M_Z^2 \end{aligned}$$

$$+ \frac{1}{22} \frac{t^2 + 1}{t^2 - 1} S_0 \left(1 - \frac{\alpha_1(Q)}{\alpha_G} \right) \quad (20)$$

where $C_\mu = \frac{1}{2} D_0 (1 - D_0) (H_3/F)^2 + e - g/t^2$, $t \equiv \tan\beta$, and $D_0 \cong 1 - m_t^2/(200 \sin\beta)^2$. D_0 vanishes at the t-quark Landau pole (for $m_t = 175$ GeV, $D_0 \leq 0.23$) and $A_R = A_t - m_{1/2}(H_2 - H_3/F)$ is the residue at the Landau pole ($A_R \cong A_t - 0.61(\alpha_3/\alpha_G)m_{1/2}$), i.e. $A_0 = A_R/D_0 - (H_3/F)m_{1/2}$ where A_0 is the cubic soft SUSY breaking parameter at M_G ²⁸. α_G is the GUT coupling constant ($\alpha_G \simeq 1/24$) and

$$S_0 = \text{Tr}(Y m^2) = (m_{H_2}^2 - m_{H_1}^2) + \sum_{i=1}^{ng} (m_{q_L}^2 - 2m_{u_R}^2 + m_{d_R}^2 - m_{\ell_L}^2 + m_{e_R}^2) \quad (21)$$

In Eq.(20), the form factors e.g. H_2 , H_3 , F are given in Ibañez et al¹⁶, n_g is the number of generations, and the (mass)² in Eq.(21) are all to be taken at M_G . We note that S_0 vanishes for universal soft breaking masses, and reduces to $S_0 = m_{H_2}^2 - m_{H_1}^2$ for three generation GUT models obeying Eq.(19). (The coefficient $(1 - \alpha_1(M_Z)/\alpha_G)/22 \cong 0.0268$ and so this term is generally small.)

The renormalization group equations evolve the universal gaugino mass $m_{1/2}$ at M_G to separate masses for SU(3), SU(2) and U(1) at M_Z :

$$\tilde{m}_i = (\alpha_i(M_Z)/\alpha_G)m_{1/2}; \quad i = 1, 2, 3 \quad (22)$$

where at 1-loop, the gluino mass $m_{\tilde{g}} = \tilde{m}_3$ ²⁹. The breaking of SU(2) x U(1) causes a mixing of the gaugino-higgsino states giving rise to two charginos $\chi_i^\pm$, $i=1,2$ and four neutralinos χ_i^0 , $i = 1 \cdots 4$. In the $\tilde{W}^\pm - \tilde{H}^\pm$ basis, the chargino mass matrix is

$$M_\chi^\pm = \begin{pmatrix} \tilde{m}_2 & \sqrt{2}M_W \sin\beta \\ \sqrt{2}M_W \cos\beta & \mu \end{pmatrix} \quad (23)$$

and in the $(\tilde{W}_3, \tilde{B}, \tilde{H}_1, \tilde{H}_2)$ basis and the neutralino mass matrix is

$$M_\chi^0 = \begin{pmatrix} \tilde{m}_2 & o & a & b \\ o & \tilde{m}_1 & c & d \\ a & c & o & -\mu \\ b & d & -\mu & o \end{pmatrix} \quad (24)$$

where $a = M_Z \cos\theta_W \cos\beta$, $c = -\tan\theta_W a$, $b = -M_Z \cos\theta_W \sin\beta$, $d = -\tan\theta_W b$ and θ_W is the weak mixing angle. In the domain where $\mu^2/M_Z^2 \gg 1$ one has

$$\begin{aligned}
2m_{\chi_1}^0 &\cong m_{\chi_1}^\pm \cong m_{\chi_2}^0 \simeq \frac{1}{3}m_{\tilde{g}} \\
m_{\chi_3}^0 &\cong m_{\chi_4}^0 \cong m_{\chi_2}^\pm \simeq \mu \gg m_{\chi_1}^0
\end{aligned} \tag{25}$$

i.e. the light states are mostly gauginos and the heavy states mostly higgsinos.

The large top mass also causes L-R mixing in the stop mass matrix

$$M_t^2 = \begin{pmatrix} m_{t_L}^2 & -m_t(A_t + \mu \tan\beta) \\ -m_t(A_t + \mu \tan\beta) & m_{t_R}^2 \end{pmatrix} \tag{26}$$

where

$$m_{t_L}^2 = m_{Q_L}^2 + m_t^2 + \left(\frac{1}{2} - \frac{2}{3} \sin^2\theta_W \right) M_Z^2 \cos 2\beta \tag{27}$$

$$m_{t_R}^2 = m_U^2 + m_t^2 + \frac{2}{3} \sin^2\theta_W M_Z^2 \cos 2\beta \tag{28}$$

and

$$\begin{aligned}
m_{Q_L}^2 &= \left[\left(\frac{1+D_0}{2} \right) + \frac{5+D_0}{6} \delta_3 - \frac{1-D_0}{6} (\delta_2 + \delta_4) \right] m_0^2 \\
&- \frac{1}{6} (1-D_0) \frac{A_R^2}{D_0} + C_Q m_{1/2}^2 - \frac{1}{66} S_0 (1 - \alpha_1(Q)/\alpha_G)
\end{aligned} \tag{29}$$

$$\begin{aligned}
m_U^2 &= \left[D_0 + \frac{2+D_0}{3} \delta_4 - \frac{1-D_0}{6} (\delta_2 + \delta_3) \right] m_0^2 \\
&- \frac{1}{3} (1-D_0) \frac{A_R^2}{D_0} + C_U m_{1/2}^2 + (2/33) S_0 (1 - \alpha_1(Q)/\alpha_G)
\end{aligned} \tag{30}$$

The coefficients C_Q , C_U are given by

$$C_Q = \frac{1}{3} \left[-\frac{1}{2} D_0 (1-D_0) (H_3/F)^2 + e \right] + \frac{\alpha_G}{4\pi} \left(\frac{8}{3} f_3 + f_2 - \frac{1}{15} f_1 \right) \tag{31}$$

$$C_U = \frac{2}{3} \left[-\frac{1}{2} D_0 (1-D_0) (H_3/F)^2 + e \right] + \frac{\alpha_G}{4\pi} \left(\frac{8}{3} f_3 - f_2 + \frac{1}{3} f_1 \right) \tag{32}$$

the form factors f_i being given in Ibañez et al.¹⁶.

The simplest model is the one with universal soft breaking masses, i.e. $\delta_i=0$. This model depends on only the four SUSY parameters and one sign

$$m_0, \quad m_{1/2}, \quad A_0, \quad B_0, \quad \text{sign}(\mu) \tag{33}$$

Alternately one may choose m_0 , $m_{\tilde{g}}$, A_t , $\tan\beta$, and $\text{sign}(\mu)$ as the independent parameters. This case has been extensively discussed in the literature³⁰.

As can be seen above, however, the deviations from universality can significantly effect the values of μ^2 and the stop masses, and these parameters play a crucial role in predictions of the theory. Thus more recently, efforts to include non-universalities into calculations have been made, and some of these effects will be discussed in the context of dark matter in Sec.4.

4. Phenomenological Implications of Supergravity Models

We shall discuss the implications of supergravity models mostly in the context of the parameter space of Eq.(33) although many of the results of the analysis are valid for a broader class of supergravity models which include non-universalities as discussed in Secs.2 and 3. We shall analyse the effect of non-universalities discussed in Sec.3 in some detail in the context of dark matter detectors. There are many phenomenological implications of supergravity unified models. One of the important implications concerns the signatures of supersymmetry in collider experiments. Such signatures depend crucially on the stability of the lowest supersymmetric particle (LSP). A stable LSP can be achieved by the imposition of an R parity symmetry. This is an attractive possibility as it also provides a candidate for cold dark matter. In fact using the renormalization group analysis discussed earlier, one finds that over most of the parameter space of supergravity unified models the lightest neutralino is the LSP³¹. In this circumstance the decay of the supersymmetric particles will have missing energy signals and at least one of the carriers of missing energy will be the neutralino. Signals of this type were studied early on after the advent of supergravity models in the supersymmetric decays of the W and Z bosons³² and such analyses have since been extended to the decays of all of the supersymmetric particles. Using these decay patterns one finds a variety of supersymmetric signals for SUSY particles at colliders where SUSY particles are expected to be pair produced when sufficient energies in the center of mass system are achieved. One signal of special interest in the search for supersymmetry is the trileptonic signal³³. For example in $p\bar{p}$ collisions one can have $p\bar{p} \rightarrow \tilde{\chi}_1^\pm + \tilde{\chi}_2^0 + X \rightarrow (l_1\bar{\nu}_1\tilde{\chi}_1^0) + (l_2\bar{l}_2\tilde{\chi}_1^0) + X$ which gives a signal of three leptons and missing energy. Using this signal a chargino of mass up to 230 GeV can be explored at the Tevatron with an integrated luminosity of³⁴ $10fb^{-1}$.

One of the most accurate probes of new physics beyond the Standard Model (SM) is provided by (g-2) for the muon and for the electron³⁵. Currently the experimental value of $a_\mu \equiv (g_\mu/2-1)$ is $a_\mu^{exp} = 11659230(84) \times 10^{-10}$ where as $a_\mu^{theory}(SM) = 11659172(15.4) \times 10^{-10}$. Here $a_\mu^{theory}(SM)$ contains the $O(\alpha^5)$ Q.E.D. contributions, $O(\alpha^2)$ hadronic corrections and up to two loop

corrections from the SM electro-weak sector. Essentially all of the error in the theoretical computations comes from the hadronic contributions which give $a_\mu^{hadron}(SM) = 687.0(15.4) \times 10^{-10}$ while the contribution from the electro-weak sector has a very small error³⁶, i.e., $a_\mu^{EW}(SM) = 15.1(0.4) \times 10^{-10}$. It is expected that the new experiment underway at Brookhaven will reduce the experimental uncertainty by a factor of about 20. Further, a new analysis by Alemany et.al.³⁷ using the τ data at LEP indicates a significant reduction in the hadronic error, i.e., $a_\mu^{hadron}(SM) = 701.4(9.4) \times 10^{-10}$. The hadronic error is expected to reduce further perhaps by another factor of two when new data from VEPP-2M, DAΦNE, and BEPC experiments comes in. Thus the improved determination of $g_\mu - 2$ in the Brookhaven experiment as well as the expected reduction in the hadronic error will allow one to test the electro-weak SM contribution.

It was pointed out in ref.³⁸ that any experiment which tests the SM electro-weak contribution to $g_\mu - 2$ can also test the supersymmetric contribution. This conclusion is supported by the recent precision analyses of $g_\mu - 2$ using the renormalization group^{39,40}. Thus the Brookhaven experiment is an important probe of supergravity grand unification.

Flavor changing neutral current processes also provide an important constraint on supergravity unified models. A process of great interest here is the decay $b \rightarrow s + \gamma$ which has been observed by the CLEO collaboration⁴¹ with a branching ratio of $BR(b \rightarrow s + \gamma) = (2.32 \pm 0.67) \times 10^{-4}$. In the SM this decay proceeds at the loop level with the exchange of W and Z bosons and the most recent analyses⁴² of the branching ratio including leading and most of the next to leading order QCD corrections yield $BR(b \rightarrow s + \gamma) = (3.48 \pm 0.31) \times 10^{-4}$ for $m_t = 176$ GeV. In supersymmetry there are additional diagrams which contribute to this process⁴³. Thus in SUGRA unification one has contributions from the exchange of the charged Higgs, the charginos, the neutralinos and from the gluino. Interestingly one finds that while the contribution from the charged Higgs exchange is always positive⁴⁴ over much of the parameter space its effects are small since $H^\pm$ is heavy. The contribution from the exchange of the other SUSY particles can be either positive or negative with the contribution of the charginos being usually the dominant one⁴⁵. Thus supergravity can accomodate a value of the $BR(b \rightarrow s + \gamma)$ which is lower than the SM value^{45,46}. If the more accurate experimental determination of $BR(b \rightarrow s + \gamma)$ continues to show a trend with a value of $BR(b \rightarrow s + \gamma)$ lower than what the SM predicts, the results would provide a strong hint for a low lying chargino and a low lying third generation squark. The $b \rightarrow s + \gamma$ experiment also imposes an important constraint on dark matter analyses and we discuss this below.

As mentioned earlier one of the remarkable results of supergravity grand

unification with R parity invariance is the prediction that the lightest neutralino χ_1^0 is the LSP over most of the parameter space. In this part of the parameter space the χ_1^0 is a candidate for cold dark matter (CDM). We discuss now the relic density of χ_1 within the framework of the Big Bang Cosmology. The quantity that is computed theoretically is $\Omega_{\chi_1} h^2$ where $\Omega_{\chi_1} = \rho_{\chi_1}/\rho_c$, ρ_{χ_1} is the neutralino relic density and ρ_c is the critical relic density needed to close the universe, $\rho_c = 3H^2/8\pi G_N$, and $H = h100km/sMpc$ is the Hubble constant. The most conservative constraint on $\Omega_{\chi_1} h^2$ is $\Omega_{\chi_1} h^2 < 1$. A variety of more constraining possibilities have also been discussed in the literature. One of the important elements in the computation of the relic density concerns the correct thermal averaging of the quantity $\langle\sigma v\rangle$ where σ is the neutralino annihilation cross section in the early universe and v is the relative neutralino velocity. Normally the thermal average is calculated by first making the approximation $\sigma v = a + bv^2$ and then evaluating its thermal average⁴⁷. However, it is known that such an approximation breaks down in the vicinity of thresholds and poles⁴⁸. Precisely such a situation exists for the case of the annihilation of the neutralino through the Z and Higgs poles. An accurate analysis of the neutralino relic density in the presence of Z and Higgs poles was given in ref.⁴⁹ and similar analyses have also been carried out since by other authors^{50,51}.

There are a number of possibilities for the detection of dark matter both direct and indirect^{52,51}. We discuss here the direct method which involves the scattering of incident neutralino dark matter in the Milky Way from nuclei in terrestrial targets. The event rates consist of two parts⁵³: one involves an axial interaction and the other a scalar interaction. The axial (spin dependent) part R_{SD} falls off as $R_{SD} \sim 1/M_N$ for large M_N where M_N is the mass of the target nucleus, while the scalar (spin independent) part behaves as $R_{SI} \sim M_N$ and increases with M_N . Thus for heavy target nuclei the spin independent part R_{SI} dominates over most of the parameter space of the model. Analysis of event rates for the scattering of neutralinos off targets such as Ge and Xe indicates that the event rates can lie over a wide range, from O(1) event/kg d to O($10^{-(5-6)}$) event/kg d⁵⁴. Inclusion of non-universalities is seen to produce definite signatures in the event rate analysis^{27,55}. Thus the minimum event rates in the region $m_{\chi_1} < 65$ GeV can be enhanced (for the case $\delta_1 = -1 = -\delta_2$) or reduced (for the case $\delta_1 = 1 = -\delta_2$) by a factor of O(10-100) due to the presence of non-universalities²⁷, while the effect of non-universalities is generally small for $m_{\chi_1} > 65$ GeV. To illustrate how this comes about we see that from Eq.(20) the non-universality correction to μ is

given by

$$\Delta\mu^2 = \frac{t^2}{t^2-1} \left[\left(\frac{1-D_0}{2}(\delta_3+\delta_4) - \frac{1+D_0}{2}\delta_2 + \frac{1}{t^2}\delta_1 \right) \right] m_0^2 \quad (34)$$

μ^2 influences detection event rates in that an increase (decrease) in μ^2 reduces (increases) R_{SI} . Thus for $\delta_1 = -1 = -\delta_2$ one finds that $\Delta\mu^2$ is negative and tends to raise R_{SI} . In addition, this could drive μ^2 negative, eliminating such points from the parameter space of the theory. These effects are more significant for small $\tan\beta$ values and thus drive the minimum event rates higher in the region $m_{\chi_1} < 65$ GeV. For $\delta_1 = -1 = -\delta_2$ one has $\Delta\mu^2 > 0$ and the effect here is opposite to that for the previous case. (Note also since D_0 is small, δ_3 and δ_4 can cancel or enhance the effects of δ_1 and δ_2). Further, it is found that the event rates for $\mu < 0$ are much suppressed (by a factor ≈ 100) compared to the case $\mu > 0$ ⁵⁴. This feature persists in the presence of non-universalities²⁷.

The detection sensitivity of current detectors is typically of order of a few events/kg d⁵⁶ which is as yet insufficient to probe a significant part of the parameter space of SUGRA models. One needs more sensitive detectors with sensitivities 2-3 orders of magnitude better to sample a significant part of the parameter space of SUGRA models. (See in this context ref.⁵⁷).

As mentioned already the minimal supergravity grand unification gives a result for $\alpha_3(M_Z)$ which is about 2-3 std larger than the current world average^{14,15}. However, because of the proximity of the GUT scale to the Planck scale one can expect corrections of size $O(M_G/M_{Pl})$ where M_{Pl} is the Planck mass. For example, Planck scale corrections can modify the gauge kinetic energy function so that one has for the gauge kinetic energy term $-(1/4)f_{\alpha\beta}F_{\alpha}^{\mu\nu}F_{\beta\mu\nu}$. For the minimal SU(5) theory, Eq.(3) can give terms $f_{\alpha\beta} = \delta_{\alpha\beta} + (c/2M_{Pl})d_{\alpha\beta\gamma}\Sigma^{\gamma}$ where Σ is the scalar field in the 24 plet of SU(5). After the spontaneous breaking of SU(5) and a re-diagonalization of the gauge kinetic energy function, one finds a splitting of the $SU(3) \times SU(2) \times U(1)$ gauge coupling constants at the GUT scale. These splittings can easily generate the 2 std correction to $\alpha_s(M_Z)$. In fact, using the LEP data one can put constraints on c . One finds that¹⁵ $-1 \leq c \leq 3$. The Planck scale correction also helps relax the stringent constraint on $\tan\beta$ imposed by $b - \tau$ unification. Thus in the absence of Planck scale correction one has that $b - \tau$ unification requires $\tan\beta$ to lie in two rather sharply defined corridors⁵⁸. One of these corresponds to a small value of $\tan\beta$, i.e., $\tan\beta \sim 2$ and the second a large value $\tan\beta \sim 50$. This stringent constraint is somewhat relaxed by the inclusion of Planck scale corrections¹⁵.

SUSY grand unified models contain many sources of proton instability. Thus in addition to the p decay occuring via the exchange of superheavy vector

lepto-quarks, one has the possibility of p decay from dimension (dim) 4 (dim 3 in the superpotential) and dim 5 (dim 4 in the superpotential) operators⁵⁹. The lepto-quarks exchange would produce $p \rightarrow e^+\pi^0$ as its dominant mode with an expected lifetime⁶⁰ of $\sim 1 \times 10^{35 \pm 1} [M_X/10^{16}]^4 y$ where for $SU(5)$ $M_X \cong 1.1 \times 10^{16}$ while the sensitivity of Super Kamiokande (Super-K) to this mode^{61,62} is 1×10^{34} . Thus the $e^+\pi^0$ mode may be at the edge of being accessible at Super-K. Proton decay from dim 4 operators is much too rapid but is easily forbidden by the imposition of R parity invariance. The p decay from dim 5 operators is more involved. It depends on both the GUT physics as well as on the low energy physics such as the masses of the squarks and of the gauginos. Analysis in supergravity unified models^{63–65} shows that one can make concrete predictions of the p decay modes within these models. Thus one can show that with the Higgs triplet mass $M_{H_3} < 10M_G$, the parameter space of minimal $SU(5)$ SUGRA will come close to being exhausted for $m_0 < 1TeV$ and $m_{\tilde{g}} < 1TeV$ if Super-K⁶² and Icarus⁶⁶ reach the sensitivity for $\bar{\nu}K^+$ mode of $2 \times 10^{34} y$ ^{64,65}.

Precision determination of soft SUSY breaking parameters can be utilized as a vehicle for the test of the predictions of supergravity grand unification. It was recently proposed that precision measurement of the soft breaking parameters can also act as a test of physics at the post GUT and string scales⁶⁷. Thus, for example, if one has a concrete model of the soft breaking parameters at the string scale then these parameters can be evolved down to the grand unification scale leading to a predicted set of non-universalities there. If the SUSY particle spectra and their interactions are known with precision at the electro-weak scale, then this data can be utilized to test a specific model at the post GUT or string scales. Future colliders such as the LHC⁶⁸ and the NLC⁶⁹ will allow one to make mass measurements with significant accuracy. Thus accuracies of up to a few percent in the mass measurements will be possible at these colliders allowing a test of post GUT and string physics up to an accuracy of $\sim 10\%$ ⁶⁷.

5. Conclusion

Supergravity grand unification provides a framework for the supersymmetric unification of the electro-weak and the strong interactions where supersymmetry is broken spontaneously by a super Higgs effect in the hidden sector and the breaking communicated to the visible sector via gravitational interactions. The minimal version of the model based on a generation independent Kahler potential contains only four additional arbitrary parameters and one sign in terms of which all the SUSY mass spectrum and all the SUSY interaction

structure is determined. This model is thus very predictive. A brief summary of the predictions and the phenomenological implications of the model was given. Many of the predictions of the model can be tested at current collider energies and at energies that would be achievable at future colliders. Other predictions of the model can be tested in non-accelerator experiments such as at Super Kamiokande and at Icarus, and by dark matter detectors. We also discussed here extensions of the minimal supergravity model to include non-universalities in the soft SUSY breaking parameters. Some of the implications of these non-universalities on predictions of the model were discussed. Future experiments should be able to see if the predictions of supergravity unification are indeed verified in nature.

Acknowledgements: This work was supported in part by NSF grant numbers PHY-9411543 and PHY-96020274.

References

1. Yu A. Golfand and E.P. Likhtman, JETP Lett. **13**, 452 (1971); D. Volkov and V.P. Akulov, JETP Lett. **16**, 438 (1972).
2. J. Wess and B. Zumino, Nucl. Phys. **B78**, 1 (1974).
3. H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
4. E. Gildener, Phys. Rev. **D14**, 1667 (1976).
5. S. Dimopoulos and H. Georgi, Nucl. Phys. **B193**, 150 (1981).
6. S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. **D24**, 1681 (1981); N. Sakai, Z. Phys. **C11**, 153 (1981).
7. L. Giradello and M.T. Grisaru, Nucl. Phys. **B194**, 65 (1982).
8. For a review of the properties of the MSSM see H. Haber and G. Kane, Phys. Rep. **117**, 75 (1985).
9. D.Z. Freedman, P. van Nieuwenhuizen and S. Ferrara, Phys. Rev. **D14**, 912 (1976); S. Deser and B. Zumino, Phys. Lett. **B62**, 335 (1976).
10. A.H. Chamseddine, R. Arnowitt and P. Nath, Phys. Rev. Lett. **49**, 970 (1982).

11. For reviews see P. Nath, R. Arnowitt and A.H. Chamseddine, *Applied N =1 Supergravity* (World Scientific, Singapore, 1984); H.P. Nilles, Phys. Rep. **110**, 1 (1984); R. Arnowitt and P. Nath, Proc. of VII J.A. Swieca Summer School ed. E. Ebohi (World Scientific, Singapore, 1994).
12. Alternate possibilities are considered elsewhere in this book.
13. P. Langacker, Proc. PASCOS 90-Symposium, Eds. P. Nath and S. Reucroft (World Scientific, Singapore 1990); J. Ellis, S. Kelley and D.V. Nanopoulos, Phys. Lett. **B249**, 441 (1990); **B260**, 131 (1991); U. Amaldi, W. de Boer and H. Furstenau, Phys. Lett. **B260**, 447 (1991); F. Anselmo, L. Cifarelli, A. Peterman and A. Zichichi, Nuov. Cim. **104A**, 1817 (1991); **115A**, 581 (1992).
14. P.H. Chankowski, Z. Pluciennik, and S. Pokorski, Nucl. Phys. **B439**, 23 (1995); J. Bagger, K. Matchev and D. Pierce, Phys. Lett **B348**, 443 (1995); R.Barbieri and L.J. Hall, Phys. Rev. Lett. **68**, 752 (1992); L.J. Hall and U. Sarid, Phys. Rev. Lett. **70**, 2673 (1993); P. Langacker and N. Polonsky, Phys. Rev. **D47**, 4028 (1993).
15. T. Dasgupta, P. Mamales and P. Nath, Phys. Rev. **D52**, 5366 (1995); D. Ring, S. Urano and R. Arnowitt, Phys. Rev. **D52**, 6623 (1995); S. Urano, D. Ring and R. Arnowitt, Phys. Rev. Lett. **76**, 3663 (1996); P. Nath, Phys. Rev. Lett. **76**, 2218 (1996).
16. K. Inoue et al., Prog. Theor. Phys. **68**, 927 (1982); L. Ibañez and G.G. Ross, Phys. Lett. **B110**, 227 (1982); L. Alvarez-Gaumé, J. Polchinski and M.B. Wise, Nucl. Phys. **B221**, 495 (1983); J. Ellis, J. Hagelin, D.V. Nanopoulos and K. Tamvakis, Phys. Lett. **B125**, 2275 (1983); L. E. Ibañez and C. Lopez, Phys. Lett. **B128**, 54 (1983); Nucl. Phys. **B233**, 545 (1984); L.E. Ibañez, C. Lopez and C. Muñoz, Nucl. Phys. **B256**, 218 (1985).
17. R. Barbieri, S. Ferrara and C.A. Savoy, Phys. Lett. **B119**, 343 (1982).
18. L. Hall, J. Lykken and S. Weinberg, Phys. Rev. **D27**, 2359 (1983).
19. P. Nath, R. Arnowitt and A.H. Chamseddine, Nucl. Phys. **B227**, 121 (1983).
20. S. Soni and A. Weldon, Phys. Lett. **B126**, 215 (1983).
21. V.S. Kaplunovsky and J. Louis, Phys. Lett. **B306**, 268 (1993).

22. E. Cremmer, S. Ferrara, L. Girardello and A. van Proeyen, Phys. Lett. **116B**, 231(1982).
23. C.T. Hill, Phys. Lett. **B135**, 47 (1984); Q. Shafi and C. Wetterich, Phys. Rev. Lett. **52**, 875 (1984).
24. SO(10) models exist with an intermediate breaking scale. However, such models may have additional difficulties in satisfying grand unification and proton decay constraints²⁵.
25. S. Urano and R. Arnowitt, hep-ph/9611389.
26. G. Gamberini, G. Ridolfi and F. Zwirner, Nucl. Phys. **B331**, 331 (1990); R. Arnowitt and P. Nath, Phys. Rev. **D46**, 3981 (1992). as this
27. P. Nath and R. Arnowitt, hep-ph/9701301, (to be pub. Phys.Rev.D)
28. P. Nath, J. Wu and R. Arnowitt, Phys. Rev. **D52**, 4169 (1995).
29. QCD corrections to this result are discussed in S.P. Martin and M.T. Vaughn, Phys. Lett. **B318**, 331 (1993); D. Pierce and A. Papadopoulos, Nucl. Phys. **B430**, 278 (1994).
30. G. Ross and R.G. Roberts, Nucl. Phys. **B377**, 571 (1992); R. Arnowitt and P. Nath, Phys. Rev. Lett. **69**, 725 (1992); M. Drees and M.M. Nojiri, Nucl. Phys. **B369**, 54 (1993); S. Kelley *et. al.*, Nucl. Phys. **B398**, 3 (1993); M. Olechowski and S. Pokorski, Nucl. Phys. **B404**, 590 (1993); G. Kane, C. Kolda, L. Roszkowski and J. Wells, Phys. Rev. **D49**, 6173 (1994); D.J. Castaño, E. Piard and P. Ramond, Phys. Rev. **D49**, 4882 (1994); W. de Boer, R. Ehret and D. Kazakov, Z.Phys. **C67**, 647 (1995); V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D49**, 4908 (1994); H. Baer, M. Drees, C. Kao, M. Nojiri and X. Tata, Phys. Rev. **D50**, 2148 (1994); H. Baer, C.-H. Chen, R. Munroe, F. Paige and X. Tata, Phys. Rev. **D51**, 1046 (1995).
31. R. Arnowitt and P. Nath, Phys. Rev. Lett. **69**, 725(1992); P. Nath and R. Arnowitt, Phys. Lett. **B289**, 368(1992).
32. S. Weinberg, Phys. Rev. Lett. **50**, 387(1983); R. Arnowitt, A.H. Chamseddine and P. Nath, Phys. Rev. Lett. **50**, 232(1983); Phys. Lett. **129B**, 445(1983); D.A. Dicus, S. Nandi, W.W. Repko and X. Tata, Phys. Lett. **129B**, 451(1983); J. Ellis, J.S. Hagelin, D.V. Nanopoulos, and M. Srednicki, Phys. Rev. Lett. **127B**, 233(1983).

33. P. Nath and R. Arnowitt, Mod. Phys. Lett. **A2**, 331(1987); R. Barbieri, F. Caravaglio, M. Frigeni, and M. Mangano, Nucl. Phys. **B367**, 28(1991); H. Baer and X. Tata, Phys. Rev. **D47**, 2739(1992); J.L. Lopez, D.V. Nanopoulos, X. Wang and A. Zichichi, Phys. Rev. **D48**, 2062(1993); H. Baer, C. Kao and X. Tata, Phys. Rev. **D48**, 5175(1993).
34. D. Amidie and R. Brock, "Report of the tev-2000 Study Group", FERMILAB-PUB-96/082.
35. T. Kinoshita and W. J. Marciano, in *Quantum Electrodynamics*, edited by T. Kinoshita (World Scientific, Singapore, 1990).pp. 419-478.
36. A. Czarnecki, B. Krauss and W. Marciano, Phys.Rev. **D52**, 2619(1995)
37. R. Alemany, M. Davier and A. Hocker, hep-ph/9703220.
38. T. C. Yuan, R. Arnowitt, A.H. Chamseddine and P. Nath, Z. Phys. **C26**, 407(1984); D. A. Kosower, L. M. Krauss, N. Sakai, Phys. Lett. **133B**, 305(1983).
39. J. Lopez, D.V. Nanopoulos, and X. Wang, Phys. Rev. **D49**, 366(1994).
40. U. Chattopadhyay and P. Nath, Phys. Rev. **D53**, 1648(1996); M. Carena, G.F. Giudice and C.E.M. Wagner, CERN-TH/96-271.
41. M. S. Alam et. al. (CLEO Collaboration), Phys. Rev. Lett.**74**,2885(1995).
42. A. J. Buras, M. Misiak, M. Munz and S. Pokorski, Nucl. Phys. **B424**, 374(1994); M. Ciuchini et. al., Phys. Lett. **B316**, 127(1993); K. Chetyrkin, M. Misiak, and M. Munz, Phys.Lett. **B400**, 206 (1997); A. J. Buras, A. Kwiatkowski and N. Pott, hep-ph/9707482.
43. S. Bertolini, F. Borzumati and A. Masiero, Phys. Rev. Lett. **59**, 180(1987); R. Barbieri and G. Giudice, Phys. Lett. **B309**, 86(1993).
44. J.L. Hewett, Phys. Rev. Lett. **70**, 1045(1993); V. Barger, M. Berger, P. Ohmann. and R.J.N. Phillips, Phys. Rev. Lett. **70**, 1368(1993).
45. M. Diaz, Phys. Lett. **B304**, 278(1993); J. Lopez, D.V. Nanopoulos, and G. Park, Phys. Rev. **D48**, 974(1993); R. Garisto and J.N. Ng, Phys. Lett. **B315**, 372(1993); J. Wu, R. Arnowitt and P. Nath, Phys. Rev. **D51**, 1371(1995); V. Barger, M. Berger, P. Ohman and R.J.N. Phillips, Phys. Rev. **D51**, 2438(1995); H. Baer and M. Brhlick, hep-ph/9610224.

46. P. Nath and R. Arnowitt, Phys. Lett. **B336**, 395(1994); F. Borzumati, M. Drees, and M.M. Nojiri, Phys. Rev.**D51**, 341(1995).
47. For a review see, E. W. Kolb and M. S. Turner, *The Early Universe* (Addison- Wesley, Redwood City, CA, 1989).
48. K. Greist and D. Seckel, Phys. Rev. **D43**, 3191 (1991); P. Gondolo and G. Gelmini, Nucl. Phys. **B360**, 145(1991).
49. R. Arnowitt and P. Nath, Phys. Lett. **B299**, 58(1993); **303**, 403(E)(1993); P. Nath and R. Arnowitt, Phys. Rev. **70**, 3696(1993).
50. H. Baer and M. Brhlick, Phys. Rev. **D53**, 597(1996); V. Barger and C. Kao, hep-ph/970443.
51. For a review see, P. Nath and R. Arnowitt, "Supersymmetric Dark Matter", in proceedings of the Heidelberg Workshop on "Aspects of Dark Matter in Astro- and Particle Physics", Heidelberg, September 16-20,1996.
52. For a review see, G. Jungman, M. Kamionkowski and K. Greist, Phys. Rep. **267**,195(1995).
53. M.W. Goodman and E. Witten, Phys. Rev. **D31**, 3059(1983); K.Greist, Phys. Rev. **D38**, (1988)2357; **D39**,3802(1989)(E); J.Ellis and R.Flores, Phys. Lett. **B300**,175(1993); R. Barbieri, M. Frigeni and G.F. Giudice, Nucl. Phys. **B313**,725(1989); M.Srednicki and R.Watkins, Phys. Lett. **B225**,140(1989); R.Flores, K.Olive and M.Srednicki, Phys. Lett. **B237**,72(1990); A. Bottino etal, Astro. Part. Phys. **1**, 61 (1992); **2**, 77 (1994); V.A. Bednyakov, H. V. Klapdor-Kleingrothaus and S.G. Kovalenko, Phys. Rev. **D50**,7128(1994).
54. R.Arnawitt and P.Nath, Mod. Phys. Lett. **A 10**,1257(1995); P.Nath and R.Arnawitt, Phys. Rev. Lett.**74**,4592(1995); R.Arnawitt and P.Nath, Phys. Rev. **D54**,2374(1996); E.Diehl, G.Kane, C.Kolda and J.Wells, Phys. Rev.**D52**,4223(1995); V. A. Bednyakov, S.G. Kovalenko, H.V. Klapdor-Kleingrothaus, and Y. Ramachers, Z. Phys. **A357**, 339(1997); L. Bergstrom and P.Gondolo, Astropart.Phys.**5**, 263(1996); J.D. Vergados, J. Phys. **G22**, 253 (1996);
55. V. Berezhinsky, A. Bottino, J. Ellis, N. Forrenco, G. Mignola, and S. Scopel, Astropart.Phys.**5**, 1(1996).

56. R. Bernabei et. al., Phys. Lett. **B389**, 757(1996); A. Bottino, F. Donato, G. Mignoa, S. Scopel, P.Belli, and A. Incicchitti, Phys.Lett.**B402**, 113(1997).
57. D.Cline, Nucl.Phys.B (Proc.Suppl.) **51B**, 304 (1996); P.Benetti et al, Nucl. Inst. and Method for Particle Physics Research, **A307**,203 (1993).
58. V. Barger, M.S. Berger, and P. Ohman, Phys. Lett. **B314**, 351(1993); W. Bardeen, M. Carena, S. Pokorski, and C.E.M. Wagner, Phys. Lett. **B320**, 110(1994).
59. S.Weinberg, Phys.Rev.**D26**,287(1982); N.Sakai and T.Yanagida, Nucl.Phys.**B197**, 533(1982); S.Dimopoulos, S.Raby and F.Wilcek, Phys.Lett. **112B**, 133(1982); J.Ellis, D.V.Nanopoulos and S.Rudaz, Nucl.Phys. **B202**,43(1982); B.A.Campbell, J.Ellis and D.V.Nanopoulos, Phys.Lett.**141B**,299(1984); S.Chadha, G.D.Coughlan, M.Daniel and G.G.Ross, Phys.Lett.**149B**,47(1984).
60. W.J. Marciano, talk at SUSY-97, Philadelphia, May 1997.
61. Particle Data Group, Phys.Rev. **D50**,1173(1994).
62. Y.Totsuka, Proc. XXIV Conf. on High Energy Physics, Munich, 1988,Eds. R.Kotthaus and J.H. Kuhn (Springer Verlag, Berlin, Heidelberg,1989).
63. R.Arnowitt, A.H.Chamseddine and P.Nath, Phys.Lett. **156B**,215(1985); P.Nath, R.Arnowitt and A.H.Chamseddine, Phys.Rev.**32D**,2348(1985); J.Hisano, H.Murayama and T. Yanagida, Nucl.Phys. **B402**,46(1993).
64. R. Arnowitt and P. Nath, Phys. Rev. **D49**, 1479(1994).
65. P. Nath and R. Arnowitt, Proc. of the Workshop “Future Prospects of Baryon Instability Search in p-Decay and $n\bar{n}$ Oscillation Experiments”, Oak Ridge, Tennessee, U.S.A., March 28-30, 1996, ed: S.J. Ball and Y.A. Kamyshkov, ORNL-6910, p. 59.
66. Icarus Detector Group, Int. Symposium on Neutrino Astrophysics, Takayama. 1992.
67. R. Arnowitt and P. Nath, hep-ph/9701325 (to be pub. Phys. Rev. D).
68. H. Baer, C. Chen, F. Paige, and X. Tata, Phys. Rev. **DD52**, 2746(1995); I. Hinchliffe, F.E. Paige, M.D. Shapiro, J. Soderqvist and W. Yao, Phys. Rev. **D55**, 5520(1997).

69. T. Tsukamoto, K. Fujii, H. Murayama, M. Yamaguchi, and Y. Okada, Phys. Rev. **D**. **51**, 3153(1995); J.L. Feng, M.E. Peskin, H. Murayama, and X. Tata, Phys. Rev. **D****52**, 1418(1992); S. Kuhlman et. al., "Physics and Technology of the NLC: Snowmass 96", hep-ex/9605011.